

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/232198007>

# Honey: An Alternative Sports Gel

Article in *Strength and conditioning journal* · February 2002

CITATIONS

9

READS

2,576

5 authors, including:



**Richard Kreider**

Texas A&M University

350 PUBLICATIONS 5,985 CITATIONS

[SEE PROFILE](#)



**Christopher Rasmussen**

Texas A&M University

96 PUBLICATIONS 1,210 CITATIONS

[SEE PROFILE](#)



**Chad M Kerksick**

Lindenwood University

223 PUBLICATIONS 2,625 CITATIONS

[SEE PROFILE](#)



**Mike Greenwood**

Texas A&M University

221 PUBLICATIONS 2,573 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Cluster Training [View project](#)



Training and Nutrition in women trained in strength [View project](#)

# Honey: An Alternative Sports Gel

**Richard B. Kreider, PhD, FACSM, EPC;  
Christopher J. Rasmussen, CSCS; Stacy L. Lancaster, CSCS;  
Chad Kerksick, CSCS; and Michael Greenwood, CSCS\*D**  
**Exercise and Sport Nutrition Laboratory**  
**Department of Human Movement Sciences and Education**  
**The University of Memphis**



**Ann C. Grandjean**  
Column Editor

IT IS WELL KNOWN THAT INGESTING carbohydrates prior to and during endurance exercise can improve performance capacity. The primary means of enhancing carbohydrate availability prior to exercise has been by ingesting carbohydrate-rich snacks, whereas the ingestion of sports drinks (glucose/electrolyte solutions) has been shown to enhance carbohydrate availability during exercise. Ingesting carbohydrates and protein within 2 hours following exercise (e.g., 1.5 g/kg carbohydrate and 0.5 g/kg protein) has been reported to increase glycogen resynthesis and protein synthesis, promote a more optimal anabolic hormonal environment, and improve immune function. Theoretically, optimizing availability of carbohydrates prior to and during exercise as well as following exercise can help optimize exercise performance and recovery, which lead to greater training adaptations (5).

Although these may seem rather simple recommendations, different types and forms of carbohydrates have varying affects on the delivery of carbohydrates to the muscle and anabolic hor-

mones. In this regard, the type and form of carbohydrate ingested may affect the digestion rate, glucose release in the blood, and insulin response.

Over the last few years, researchers in the Exercise and Sport Nutrition Lab at the University of Memphis have been evaluating the effects of ingesting various types of carbohydrate gels prior to and during exercise on exercise capacity, as well as the effects of different types of carbohydrate/protein powders following exercise on recovery (2–6). The rationale has been to try to determine the optimal type of carbohydrate to ingest prior to, during, and/or following resistance and endurance exercise. An additional goal was to determine whether honey (in gel and powder form) can serve as a natural and less expensive source of carbohydrates.

Our most recent study evaluated the effects of ingesting honey on blood glucose, insulin, and cycling performance prior to and during endurance cycling. Our rationale was based on results of our initial study in this series (3, 6), which found that the carbohydrate

profile and glycemic index response of honey was nearly identical to that of a popular sports gel. Moreover, contrary to anecdotal myth, we found honey did not promote physical or psychological signs of hypoglycemia in fasted subjects (3, 5), during resistance training (1), or following resistance training (1, 2). In this study, 9 well-trained male cyclists performed three 40-mile time trials on their own racing bicycle attached to a computerized race simulator. Each race was separated by 1 week. Subjects were asked to prepare for each time trial as they would for a competitive race and to follow similar dietary intake the day before each time trial. In a double-blind, randomized, and counterbalanced manner, subjects ingested 15 g of a noncaloric flavored gel placebo, a dextrose gel, or honey with 250 mL of water prior to and every 10 miles during each time trial. The placebo and carbohydrate gels were packaged in generic foil packets for double-blind administration. Blood samples were taken prior to and every 10 miles during the race. In addition, power output, split times, heart rate, and ratings of

perceived exertion were determined throughout the time trials. Subjects were paid based on their performance during each time trial in order to encourage their best effort.

Results revealed that the subjects tolerated the gels well with no complaints of symptoms of hypoglycemia or gastrointestinal upset. Total time to perform the time trials (placebo  $131.3 \pm 3.6$  minutes, dextrose  $128.3 \pm 3.8$  minutes, and honey  $128.8 \pm 3.5$  minutes;  $P = 0.02$ ) was significantly faster when subjects ingested the dextrose and honey gels compared with the placebo. Average power (placebo  $164 \pm 11$  W, dextrose  $175 \pm 13$  W, and honey  $174 \pm 12$  W;  $P = 0.002$ ) was also significantly higher when subjects ingested the carbohydrate gels during exercise. Additionally, mean heart rate (placebo  $171 \pm 6$  b/min, dextrose  $178 \pm 7$  b/min, and honey  $177 \pm 5$  b/min;  $P = 0.08$ ) and glucose (placebo  $5.4 \pm 0.2$  mmol/L, dextrose  $5.8 \pm 0.3$  mmol/L, and honey  $6.0 \pm 0.4$  mmol/L;  $P = 0.10$ ) values tended to be higher in the groups ingesting carbohydrates during the time trials. No significant differences were observed in insulin (placebo  $5.3 \pm 0.3$  mIU/mL, dextrose  $5.1 \pm 0.2$  mIU/mL, and honey  $6.0 \pm 0.8$  mIU/mL;  $P = 0.29$ ). These findings indicate that ingesting dextrose and honey gels during endurance cycling can improve performance presumably by enhancing carbohydrate availability and work output. Moreover, honey can serve as an effective and less expensive source of carbohydrate gel. ▲

## ■ References

1. Earnest, C., R. Kreider, J. Lundberg, C. Rasmussen, P. Cowan, M. Greenwood, and A. Almada. Effects of pre-exercise carbohydrate feedings on glu-

cose and insulin responses during and following resistance exercise. *J. Strength Cond. Res.* 14:361. 2000.

2. Kreider, R., J. Lundberg, C. Rasmussen, P. Cowan, M. Greenwood, C. Earnest, and A. Almada. Effects of ingesting protein with various forms of carbohydrate following resistance-exercise on substrate availability and markers of catabolism. *J. Strength Cond. Res.* 14:366. 2000.
3. Kreider, R., C. Rasmussen, J. Lundberg, P. Cowan, M. Greenwood, C. Earnest, and A. Almada. Effects of ingesting carbohydrate gels on glucose, insulin and perception of hypoglycemia. *FASEB J.* 14: A490. 2000.
4. Lancaster, S, R.B. Kreider, C. Rasmussen, C. Kerksick, M. Greenwood, P. Milnor, A.L. Almada, and C.P. Earnest. Effects of honey supplementation on glucose, insulin and endurance cycling performance. *FASEB J.* 15:LB315. 2001.
5. Leutholtz, B., and R.B. Kreider. Optimizing nutrition for exercise and sport. In: *Nutrition Health*. T. Wilson and N.J. Temple, eds. Totowa, NJ: Humana Press, 2001. pp. 207–235.
6. Rasmussen, C., R. Kreider, J. Lundberg, P. Cowan, M. Greenwood, C. Earnest, and A. Almada. Analysis of the glycemic index and insulin response index of various carbohydrate gels. *FASEB J.* 14: A489. 2000.

Register On-line for upcoming  
Conferences and Events...

[www.nsca-lift.org](http://www.nsca-lift.org)



**All the  
RIGHT  
MOVES**



**Thank  
You...**

*to all employers  
who participated  
at **SSTC 2002** –  
San Antonio, TX*

## Employers –

**Post Your Job Vacancies  
Today FREE Of Charge!**

- Each position posted will appear on the NSCA website and in the Career Hotline.
- Complete and submit the on-line form located at [www.nsca-lift.org/CareerCenter/addlisting.htm](http://www.nsca-lift.org/CareerCenter/addlisting.htm)

**Convenient, Professional,  
User Friendly and  
Cost Effective!**

## Members Get The Latest

Check out NSCA Career Services at  
[www.nsca-lift.org](http://www.nsca-lift.org)  
for current, top-notch job listings.

## For more information:

Call 800-815-6826

E-mail [career@nsca-lift.org](mailto:career@nsca-lift.org)  
or Visit

[www.nsca-lift.org/AboutNSCA/inforequest.asp](http://www.nsca-lift.org/AboutNSCA/inforequest.asp)